

A Literature Review of Molecular Markers Predictive of Clinical Response to Cytotoxic Chemotherapy in Patients with Lung Cancer

Ikuo Sekine, MD, John D. Minna, MD,† Kazuto Nishio, MD,‡
Tomohide Tamura, MD,* and Nagahiro Saijo, MD**

Background: To find candidate genes for a predictive chemosensitivity test in patients with lung cancer by using a literature review.

Methods: Using MEDLINE searches, “in vitro chemosensitivity associated genes” and articles on association of the gene alteration with clinical chemosensitivity in lung cancer patients were selected. We calculated odds ratios (ORs) and their 95% confidence intervals (95% CIs) of response rates for patients who had tumors with or without gene alteration. Combined ORs and 95% CIs were estimated using the DerSimonian-Laird method.

Results: Of the 80 in vitro chemosensitivity-associated genes identified, 13 genes were evaluated for association with clinical chemosensitivity in 27 studies. The median (range) number of patients in each study was 50 (range, 28-108). The response rates of lung cancer with high and low P-glycoprotein expression were 0% and 73% to 85%, respectively ($p < 0.001$). Glutathione S-transferase pi expression (OR 0.22, 95% CI 0.06-0.79), excision repair cross-complementing 1 alterations (combined OR 0.53, 95% CI 0.28-1.01; $p = 0.055$), and tumor suppressor p53 mutation (combined OR 0.25, 95% CI 0.12-0.52) were associated with clinical chemosensitivity.

Conclusion: In total, 80 in vitro chemosensitivity-associated genes were identified in the literature, and high and low P-glycoprotein, glutathione S-transferase pi expression, excision repair cross-complementing 1 alterations, and tumor suppressor p53 mutation were candidates for future clinical trials of chemosensitivity tests in lung cancer patients.

Key Words: chemotherapy, drug response, molecular markers, prediction, lung cancer

(*J Thorac Oncol.* 2006;1: 31-37)

Lung cancer is the leading cause of death in many countries despite extensive basic research and clinical trials. Approximately 80% of patients with lung cancer have developed distant metastases either by the time of initial diagnosis or during recurrence after surgery for local disease. Systemic

chemotherapy against lung cancer, however, has limitations in efficacy such that patients with distant metastases rarely live long.¹

Tumor response to chemotherapy varies among patients, and objective tumor response rates to standard chemotherapy regimens are approximately 20 to 40% in patients with non-small-cell lung cancer and 60 to 90% in patients with small-cell lung cancer. Thus, it would be extremely useful to know in advance whether patients have tumors that respond to chemotherapy agents and whether the tumors would be resistant to such therapy. For this purpose, cell culture-based chemosensitivity tests have been investigated for more than 20 years, but they are not widely accepted because of technical problems such as the large amount of material required, a low success rate for the primary culture, length of time required, and poor correlation with the clinical response.²⁻⁵

To overcome these obstacles, DNA-, RNA-, and protein-based chemosensitivity tests have been created, but gene alterations that are predictive of the clinical drug response are not established. Recently, as many as 400 genes whose expression was associated with drug response were identified by cDNA microarray studies, but their functions do not seem to be related to drug sensitivity or resistance.⁶⁻¹⁰ In addition, the genes identified by microarray studies were highly unstable and depended on the selection of patients used for gene identification.^{11,12} The purpose of this study was to provide an overview of gene alterations in lung cancer that are associated with chemotherapy drug response to identify candidate genes for predictive chemosensitivity tests in patients with lung cancer.

MATERIALS AND METHODS

Because one set of genes associated with chemosensitivity is those directly involved in drug resistance mechanisms, we conducted a MEDLINE search for articles on tumor drug resistance published in the years 2001-2003. This search yielded 112 studies, including several review articles. By searching manually through these articles, we identified 134 genes or gene families that may be involved in drug resistance based on their function. We conducted the second MEDLINE searches for papers of in vitro studies on the 134 genes or gene families by using their names as a keyword.

*Divisions of Thoracic Oncology and Internal Medicine, National Cancer Center Hospital, Tokyo, Japan; †Hamon Center for Therapeutic Oncology Research, The University of Texas Southwestern Medical Center at Dallas, Dallas, TX; ‡Pharmacology Division, National Cancer Center Research Institute, Tokyo, Japan.

Copyright © 2006 by the International Association for the Study of Lung Cancer

ISSN: 1556-0864/06/0101-0031

From the 134 genes, we selected genes that met the following definition of “in vitro chemosensitivity associated genes”: 1) alteration of the gene was identified in a human drug-induced resistant, solid tumor cell line; 2) transfection of the gene induced drug resistance; or 3) down-regulation of the gene or its encode protein increased drug sensitivity. In this latter category, we included studies in which the gene expression or function was suppressed by antisense RNA, hammerhead ribozyme, or an antibody against the gene product. We excluded studies in which drugs were used to inhibit function because the specificity of the drug against the target may not have been complete. We performed a third MEDLINE search for articles on the association between the gene alteration and chemosensitivity of lung cancer cell lines by using the name of the gene as a keyword. Articles in which the association was evaluated in 20 or more cell lines were included in this study. Finally, we searched MEDLINE for studies on the association between the gene alteration and clinical drug response in patients with lung cancer by using the name of the gene as a keyword. Articles in which the association was evaluated in 25 or more patients with advanced lung cancer were included in this study. Studies in which gene expression was evaluated with microarray were excluded because result analysis and interpretation of this technique have not been established, as indicated by the fact that the list of genes identified by microarray studies was highly variable without overlap between these gene sets.^{11,12} Clinical studies on concurrent chemoradiotherapy were excluded. We constructed 2×2 tables from the response data and calculated odds ratios (ORs), their variances, and 95% confidence intervals (95% CIs) for the patients who had tumors with gene alteration relative to those who had tumors without gene alteration. Combined ORs and 95% CIs were estimated using the DerSimonian-Laird method.¹³ When a response rate was 0, association with gene alteration was evaluated using the χ^2 test because 95% CIs for ORs cannot be calculated. The name of each gene was standardized according to Human Gene Nomenclature Database of National Center for Biotechnology Information.

RESULTS

Of the 134 genes or gene families found, a gene alteration in drug-induced resistant cells, an increased or decreased resistance in transfected cells, and an altered sensitivity in gene down-regulated cells were reported for 45, 57, and 32 genes, respectively. In total, 80 genes met the definition of “in vitro chemosensitivity associated gene” (Table 1).

Gene alteration was associated with in vitro chemosensitivity in 15 (50%) of 30 studies on 15 (56%) of 27 gene alterations (Table 2). Clinical drug response was evaluated in 27 studies on 13 gene alterations. The methods used to identify gene alteration included immunohistochemical protein expression analysis ($n = 18$), polymerase chain reaction (PCR)-based mRNA expression analysis ($n = 3$), and PCR-based mutation analysis ($n = 6$). All but one clinical study was retrospective, and the median (range) number of patients in each study was 50 (28-108). Gene alteration was associated with clinical response in 8 of the 27 (30%) studies (Table 2).

TABLE 1. In Vitro Chemosensitivity-Associated Genes

Transporters: ABCA2, ABCB1, ABCB11, ABCC1, ABCC2, ABCC3, ABCC4, ABCC5, ABCG2, MVP, ATP7A, ATP7B, SLC29A1, SLC28A1, SLC19A1
Drug targets: TUBB, TUBB4, TUBA, TYMS, TOP1, TOP2A, TOP2B, DHFR,
Target-associated proteins : MAP4, MAP7, STMN1, KIF5B, HSPA5, PSMD14, FPGS
Intracellular detoxifiers: GSTP1, GPX, GCLC, GGT2, MT, RRM2, AKR1B1
DNA damage recognition and repair proteins: HMGB1, HMGB2, ERCC1, XPA, XPD, MSH2, MLH1, PMS2, APEX1, MGMT, BRCA1, GLO1
Cell cycle regulators: RB1, GML, CDKN1A, CCND1, CDKN2A, CDKN1B
Mitogenic signal regulators: ERBB2, EGFR, KRAS2, HRAS, RAF1
Survival signal regulators: AKT1, AKT2
Integrin: ITGB1
Transcription factors: JUN, FOS, MYC, NFKB1
Apoptosis regulators: TP53, MDM2, TP73, BCL2, BCL2L1, MCL1, BAX, BIRC4, BIRC5, TNFRSF6, CASP3, CASP8, HSPB1

We evaluated the association between transporter P-glycoprotein/multidrug resistance 1 (ABCB1) expression and clinical chemosensitivity in four studies. The response rate of lung cancer with high ABCB1 expression was consistently 0%, whereas that for lung cancer with low ABCB1 expression was 73 to 85% (Table 3). Among drug targets, only topoisomerase II-beta (TOP2B) expression was associated with clinical drug response in patients with small-cell lung cancer (OR 0.29, 95% CI 0.09-0.95). The intracellular detoxifier glutathione s-transferase pi (GSTP1) was associated with both in vitro and clinical drug response (OR 0.22, 95% CI 0.06-0.79) (Table 4). DNA repair gene excision repair cross-complementing 1 (ERCC1) alterations were associated with drug response among patients with non-small-cell lung cancer with marginal statistical significance; the combined OR (95% CI) for ERCC1 alteration was 0.53 (0.28-1.01; $p = 0.055$) (Table 5). Tumor suppressor p53 (TP53) mutation was the only alteration associated with drug response among patients with non-small-cell lung cancer among genes involved in cell cycle and apoptosis. A combined OR (95% CI) for TP53 among patients with non-small-cell lung cancer was 0.25 (0.12-0.52) (Table 6). B-cell CLL/lymphoma 2 (BCL2) and its family protein expression was not associated with clinical drug response (Table 7).

DISCUSSION

We identified 80 in vitro chemosensitivity-associated genes in our literature search. Of these, 13 were evaluated clinically in 27 studies; ABCB1, TOP2B, GSTP1, and ERCC1 expression and TP53 mutation were associated with changes to drug responses among patients with lung cancer.

Classical drug resistance is believed to be the result of molecular changes inhibiting the drug-target interaction. ABCB1, an ATP-binding cassette protein, acts as an energy-dependent transmembrane efflux pump and decreases the intracellular accumulation of anticancer drugs, including anthracyclines, vinca alkaloids, taxanes, and epipodophyllotox-

TABLE 2. Chemosensitivity-Associated Genes and Association with Chemosensitivity

Category	No of Genes	Association with chemosensitivity					
		In vitro studies (n)			Clinical studies (n)		
		Total	Yes	%	Total	Yes	%
Transporter	15	9	5	55	4	4	100
Drug target	8	2	1	50	5	1	20
Target-associated protein	7	0	0		0	0	
Intracellular detoxifier	7	3	3	100	1	1	100
DNA repair	10	1	1	100	6	0	0
Damage recognition protein	2	0	0		0	0	
Cell cycle	6	4	2	50	2	0	0
Mitogenic signal	5	3	1	33	1	0	0
Survival signal	2	0	0		0	0	
Transcription factor	4	3	0	0	0	0	
Cell adhesion-mediated drug-resistance protein	1	0	0		0	0	
Apoptosis	13	5	2	40	8	2	25
Total	80	30	15	50	27	8	30

TABLE 3. ABCB1 (P-Glycoprotein) and Clinical Response to Chemotherapy

Author	Histology	Drugs	Method	Expression	Patients (n)	RR (%)	Odds ratio
Yeh et al. ³⁰	Non-small cell	Paclitaxel	IHC	Low	35	80	0
				High	15	0	$p < 0.001^*$
Kawasaki et al. ³¹	Small cell	CAV or EP	IHC	Low	26	85	0
				High	4	0	$p < 0.001^*$
Hsia et al. ³²	Small cell	EP	IHC	Low	37	73	0
				High	13	0	$p < 0.001^*$
Savaraj et al. ³³	Small cell	CAV, CEV, or EP	RT-PCR	Low	24	75	0
				High	7	0	$p < 0.001^*$

Combined odds ratio for ABCB1 expression in patients with SCLC: 0

IHC, Immunohistochemical analysis; RR, response rate; RT-PCR, reverse transcriptase-polymerase chain reaction.

*Calculated using the χ^2 test because the confidence interval cannot be calculated.

ins. Overexpression of this protein gives tumor cells a multidrug resistance phenotype in vitro, which is thought to be associated with clinical chemoresistance.¹⁴ Our review showed that the response rate of tumors with ABCB1 overexpression was 0 in all studies of lung cancer, whereas that for lung cancer tumors with low ABCB1 expression was 73 to 85% (Table 3).

There is a close relationship between drug sensitivity and quantitative and qualitative alterations of the drug's target, including tamoxifen sensitivity and estrogen receptor expression and trastuzumab response and Her-2/neu overexpression in breast cancer,¹⁵ imatinib resistance and BCR-ABL gene amplification and mutations in Philadelphia chromosome-positive leukemias,¹⁶ and imatinib response and KIT gene mutations in gastrointestinal stromal tumors.¹⁷ In all of these cases, the target molecule is a receptor or a mutated tyrosine kinase located at the entry of growth-stimulating signal transduction pathways. Recently, gefitinib, a tyrosine kinase inhibitor of the epidermal growth factor receptor (EGFR), has been developed, and two large phase II trials

showed a response rate of 18% and 12% in patients with non—small-cell lung cancer who were previously treated with conventional chemotherapy.^{18,19} Responses to the drug have been unpredictable, but mutations of the EGFR gene were identified in patients with gefitinib-responsive lung cancer.^{20,21} Furthermore, all mutations in these tumors were restricted to the activation loop of the kinase domain of EGFR, which are in distinct contrast to mutations in extracellular and regulatory domains of EGFR in glioblastoma, which are unresponsive to gefitinib.²² Thus, molecular developments of structure and function of the targets hold the promise of targeted cancer therapy. The target molecules of many anticancer cytotoxic agents have not been clearly defined; therefore, the relationship between the target molecule status and sensitivity to the agent has not been established. TOP2B expression was associated with drug response in patients with small-cell lung cancer, with a response rate of 71% for high TOP2B expression tumors versus 90% for low TOP2B expression tumors (OR 0.29, 95% CI 0.09-0.95).²³ This result, however, is in contrast with the idea that a higher

TABLE 4. Drug Targets, Intracellular Detoxifier, and Clinical Response to Chemotherapy

Author	Histology	Drugs	Method	Expression	Patients (<i>n</i>)	RR (%)	Odds ratio (95% CI)
Beta-tubulin class III							
Rosell et al. ³⁴	Non-small cell	Paclitaxel,	Real-time	Low	13	46	0.39
		Vinorelbine	PCR	High	24	25	(0.09-1.62)
Topoisomerase II-alpha							
Dingemans et al. ²³	Small cell	CEV or EP	IHC	Low	65	85	0.65
				High	23	80	(0.20-2.17)
Dingemans et al. ³⁵	Non-small cell	Platinum-based	IHC	Low	30	47	0.67
				High	8	38	(0.14-3.40)
Topoisomerase II-beta							
Dingemans et al. ²³	Small cell	CEV or EP	IHC	Low	48	90	0.29
				High	35	71	(0.09-0.95)
Dingemans et al. ³⁵	Non-small cell	Platinum-based	IHC	Low	18	50	0.86
				High	13	46	(0.21-3.58)
Glutathione s-transferase pi							
Nakanishi et al. ³⁶	Non-small cell	Cisplatin-based	IHC	Low	17	47	0.22
				High	37	16	(0.06-0.79)

CI, confidence interval; IHC, immunohistochemical analysis; PCR, polymerase chain reaction; RR, response rate; CEV, cyclophosphamide, etoposide, and vincristine; EP, etoposide and cisplatin.

TABLE 5. DNA Repair Genes and Clinical Response to Chemotherapy

Author	Histology	Drugs	Method	Alteration	Patients (n)	RR (%)	Odd ratio (95% CI)
Excision repair cross-complementing 1 expression							
Lord et al. ³⁷	Non-small cell	Cisplatin, gemcitabine	Real-time PCR	Low	23	52	0.38
				High	24	36	(0.11-1.26)
Excision repair cross-complementing 1 (ERCC1) polymorphism at codon 118							
Ryu et al. ³⁸	Non-small cell	Cisplatin-based	PCR	C/C	54	54	0.61
			Hybridization	C/T or T/T	53	42	(0.28-1.31)
Combined odds ratio (95% C.I.) for ERCC1 alteration in patients with NSCLC:0.53 (0.28-1.01, <i>p</i> = 0.055)							
Xeroderma pigmentosum group D polymorphism							
At codon 231							
Ryu et al. ³⁸	Non-small cell	Cisplatin-based	PCR	G/G	100	48	1.08
			Hybridization	G/A or A/A	8	50	(0.26-4.57)
At codon 312							
Camps et al. ³⁹	Non-small cell	Cisplatin, gemcitabine	PCR	G/G	18	17	3.33
			Sequencing	G/A or A/A	15	40	(0.66-16.7)
At codon 751							
Camps et al. ³⁹	Non-small cell	Cisplatin, gemcitabine	PCR	A/A	22	23	2.04
			Sequencing	A/C or C/C	16	38	(0.49-8.45)
Ryu et al. ³⁸	Non-small cell	Cisplatin-based	PCR	A/A	96	49	0.74
			Hybridization	A/C	12	42	(0.22-2.51)
Combined odds ratio (95% CI) for XPD polymorphism in patients with NSCLC: 1.38 (0.68-2.78).							

CI, confidence interval; PCR, polymerase chain reaction; RR, response rate; NSCLC, non—small-cell lung cancer; XPD, xeroderma pigmentosum group D.

TABLE 6. Cell Cycle Regulators, Mitogenic Signals, Tumor Protein p53, and Clinical Response to Chemotherapy

Author	Histology	Drugs	Method	Alteration	Patients (n)	RR (%)	Odds ratio (95% CI)
Retinoblastoma 1 expression Gregorc et al. ⁴⁰	Non-small cell	Cisplatin-based	IHC	Low High	61 41	51 32	0.45 (0.20-1.03)
Cyclin-dependent kinase inhibitor 1A, p21 expression Dingemans et al. ²³	Small cell	CEV, EP	IHC	Low High	63 22	90 71	0.57 (0.17-1.92)
Kirsten rat sarcoma 2 viral oncogene homolog mutation Rodenhuis et al. ^{41, a}	Aenocarcinoma	Ifosfamide, carboplatin	PCR-MSH	Normal Mutated	46 16	26 19	0.65 (0.16-2.70)
Tumor protein p53 (P53) mutation Nakanishi et al. ³⁶	Non-small cell	Cisplatin-based	IHC	Normal Mutated	11 29	45 15	0.19 (0.04-0.94)
Gregorc et al. ⁴⁰	Non-small cell	Cisplatin-based	IHC	Normal Mutated	56 46	57 26	0.26 (0.11-0.62)
Combined odds ratio (95% CI) for P53 mutation in patients with NSCLC: 0.25 (0.12-0.52)							
Kawasaki et al. ³¹	Small cell	CAV or EP	IHC	Normal Mutated	10 20	70 75	1.3 (0.24-6.96)
Dingemans et al. ²³	Small cell	CEV or EP	IHC	Normal Mutated	47 45	85 82	0.81 (0.27-2.45)
Combined odds ratio (95% C.I.) for P53 mutation in patients with SCLC: 0.93 (0.37-2.35).							

CI, confidence interval; IHC, immunohistochemical analysis; PCR-MSH, polymerase chain reaction-mutation specific hybridization; RR, response rate; CEV, cyclophosphamide, etoposide, and vincristine; EP, etoposide and cisplatin.

^aProspective study.

TABLE 7. B-Cell CLL/Lymphoma 2 (BCL2) Family Expression and Clinical Response to Chemotherapy

Author	Histology	Drugs	Method	Expression	Patients (n)	RR (%)	Odds ratio (95% CI)
BCL2 Krug et al. ⁴²	Non-small cell	Docetaxel, vinorelbine	IHC	Low High	26 5	46 60	1.75 (0.25-12.3)
Dingemans et al. ²³	Small cell	CEV or EP	IHC	Low High	20 71	79 85	1.36 (0.38-4.86)
Takayama et al. ⁴³	Small cell	CAV or EP	IHC	Low High	17 21	76 62	0.50 (0.12-2.08)
Combined odds ratio (95% CI) for BCL2 expression in patients with SCLC: 0.87 (0.33-2.32)							
BAX (BCL2-associated X protein) Krug et al. ⁴²	Non-small cell	Docetaxel, vinorelbine	IHC	Low High	9 19	56 47	0.72 (0.15-3.54)

CI, confidence interval; IHC, immunohistochemical analysis; RR, response rate; CEV, cyclophosphamide, etoposide, and vincristine; EP, etoposide and cisplatin.

expression of topoisomerase II enzymes correlates with greater chemosensitivity in patients with breast cancer.²⁴

In addition to genes involved in classical drug resistance, genes that act downstream of the initial damage induced by a drug-target complex are thought to play an important role in chemosensitivity.²⁵ ERCC1 is a key enzyme in nucleotide excision repair, one of the key pathways by which cells repair platinum-induced DNA damage. High levels of ERCC1 mRNA have been associated with platinum

resistance in the treatment of ovarian and gastric cancer.^{26,27} The codon 118 in exon 4 of ERCC1 gene is polymorphic with the nucleotide alteration AAC to AAT. Although this base change results in coding for the same amino acid, it may affect gene expression based on the usage frequency of synonymous codons.²⁸ The associations between drug response and both ERCC1 gene expression and polymorphism at codon 118 in patients with non—small-cell lung cancer have been reported in the literature. A combined OR (95%

CI) for these ERCC1 alterations was 0.53 (0.28-1.01, $p = 0.055$), although each study failed to show statistical significant association. Thus, ERCC1 may be a candidate for evaluation of the predictability of drug response in future clinical trials.

TP53, which is mutated or deleted in more than half of lung cancer cells, has a remarkable number of biological activities, including cell-cycle checkpoints, DNA repair, apoptosis, senescence, and maintenance of genomic integrity. Because most anticancer cytotoxic agents induce apoptosis through either DNA damage or microtubule disruption, mutated TP53 may decrease chemosensitivity by inhibiting apoptosis or, in contrast, may increase chemosensitivity by impairing DNA repair after drug-induced DNA damage.²⁹ This review showed that mutated TP53 was associated with poor drug response in patients with non—small-cell lung cancer (Table 6).

No other genes located downstream (including xeroderma pigmentosum group D, retinoblastoma 1, cyclin-dependent kinase inhibitor 1A, Kirsten rat sarcoma 2 viral oncogene homolog, B-cell CLL/lymphoma 2, and B-cell CLL/lymphoma 2-associated X protein) were associated with clinical drug response (Tables 5-7). The association was evaluated for only 8 of 43 *in vitro* chemosensitivity-associated downstream genes; therefore, key genes may be among the remaining 35 genes. Most clinical studies included a limited number of patients with various background characteristics such as tumor stage and chemotherapy regimen administered, which resulted in low statistical power to identify the association. Finally, because all but one study was retrospective, the quality of tumor samples may vary, and it is therefore unclear whether the gene alteration was detected in all samples. Thus, in future prospective clinical studies, the method of tumor sample collection and preservation, as well as immunohistochemistry and polymerase chain reaction-based methods, should be standardized, and the sample size of patients should be determined with statistical consideration.

The recently developed microarray technique enables investigators analyze mRNA expression of more than 20,000 genes at once, and as many as 100 to 400 genes were selected statistically as chemosensitivity-related genes.^{6-8,10} Among them, however, only a limited number of genes were functionally related to chemosensitivity, and only ABCB1 and BAX corresponded with the 80 chemosensitivity-associated genes identified in this literature review, which were picked because of their known function and contribution to *in vitro* chemosensitivity. Thus, it will be interesting to evaluate the role of expression profile of these genes using microarray analysis.

The association between the expression and alterations of genes and clinical drug responses should be studied further in prospective trials. ABCB1, GSTP1, ERCC1, and TP53, and other genes identified by exploratory microarray analyses should be evaluated in those trials. Simple methods to identify gene alterations, such as immunohistochemistry and polymerase chain reaction-based techniques, will be feasible in future clinical trials because of their simplicity, cost, and

time. The median number of patients in retrospective studies analyzed in this review was 50 (range, 28-108). In future prospective trials, sample size consideration for statistical power will also be important.

In conclusion, we identified 80 *in vitro* chemosensitivity-associated genes in a review of the literature; ABCB1, GSTP1, and ERCC1 expression and TP53 mutation were associated with drug responses among patients with lung cancer.

ACKNOWLEDGMENTS

This study was supported in part by Lung Cancer SPOR Grant P50CA70907, ATP010019, and Grants-in-Aid for Cancer Research from the Ministry of Health, Labour and Welfare of Japan. We thank Yuko Yabe and Mika Nagai for their collection and arrangement of numerous papers.

REFERENCES

1. Sekine I, Saijo N. Novel combination chemotherapy in the treatment of non-small cell lung cancer. *Expert Opin Pharmacother* 2000;1:1131-1161.
2. Gazdar AF, Steinberg SM, Russell EK, et al. Correlation of *in vitro* drug-sensitivity testing results with response to chemotherapy and survival in extensive-stage small cell lung cancer: a prospective clinical trial. *J Natl Cancer Inst* 1990;82:117-124.
3. Cortazar P, Johnson BE. Review of the efficacy of individualized chemotherapy selected by *in vitro* drug sensitivity testing for patients with cancer. *J Clin Oncol*. 1999;17:1625-1631.
4. Cortazar P, Gazdar AF, Woods E, et al. Survival of patients with limited-stage small cell lung cancer treated with individualized chemotherapy selected by *in vitro* drug sensitivity testing. *Clin Cancer Res* 1997;3:741-747.
5. Shaw GL, Gazdar AF, Phelps R, et al. Individualized chemotherapy for patients with non-small cell lung cancer determined by prospective identification of neuroendocrine markers and *in vitro* drug sensitivity testing. *Cancer Res* 1993;53:5181-5187.
6. Mariadason JM, Arango D, Shi Q, et al. Gene expression profiling-based prediction of response of colon carcinoma cells to 5-fluorouracil and camptothecin. *Cancer Res* 2003;63:8791-8812.
7. Kikuchi T, Daigo Y, Katagiri T, et al. Expression profiles of non-small cell lung cancers on cDNA microarrays: identification of genes for prediction of lymph-node metastasis and sensitivity to anti-cancer drugs. *Oncogene* 2003;22:2192-2205.
8. Chang JC, Wooten EC, Tsimelzon A, et al. Gene expression profiling for the prediction of therapeutic response to docetaxel in patients with breast cancer. *Lancet* 2003;362:362-369.
9. Dan S, Tsunoda T, Kitahara O, et al. An integrated database of chemosensitivity to 55 anticancer drugs and gene expression profiles of 39 human cancer cell lines. *Cancer Res* 2002;62:1139-1147.
10. Girard L, Sekine I, Shah J, et al. Correlation between *in vitro* drug sensitivity and microarray-based gene expression signatures in lung and breast cancer. In Proceedings of the 95th Annual Meeting of American Association for Cancer Research, Orlando, FL, March 27-31, 2004. Pp. 1098.
11. Ein-Dor L, Kela I, Getz G, et al. Outcome signature genes in breast cancer: is there a unique set? *Bioinformatics* 2006;21:171-178.
12. Michiels S, Koscielny S, Hill C. Prediction of cancer outcome with microarrays: a multiple random validation strategy. *Lancet* 2006;365:488-492.
13. Armitage P, Berry G, Matthews JNS. Statistical Methods in Medical Research, 4th ed. Oxford: Blackwell Science Ltd, 2002.
14. Sekine I, Saijo N. Polymorphisms of metabolizing enzymes and transporter proteins involved in the clearance of anticancer agents. *Ann Oncol* 2001;12:1515-1525.
15. Ellis M, Hayes DF, Lippman ME. Treatment of metastatic breast cancer. In Harris J, Lippman ME, Morrow M, Osborne CK (Eds.), *Diseases of*

- the Breast, 3rd ed. Philadelphia: Lippincott Williams & Wilkins, 2003.Pp. 1101-1159.
16. Gambacorti-Passerini CB, Gunby RH, Piazza R, et al. Molecular mechanisms of resistance to imatinib in Philadelphia-chromosome-positive leukaemias. *Lancet Oncol* 2003;4:75-85.
 17. Heinrich MC, Corless CL, Demetri GD, et al. Kinase mutations and imatinib response in patients with metastatic gastrointestinal stromal tumor. *J Clin Oncol*. 2003;21:4342-4349.
 18. Fukuoka M, Yano S, Giaccone G, et al. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2003;21:2237-2246.
 19. Kris MG, Natale RB, Herbst RS, et al. Efficacy of gefitinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase, in symptomatic patients with non-small cell lung cancer: a randomized trial. *JAMA* 2003;290:2149-2158.
 20. Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004;350:2129-2139.
 21. Paez JG, Janne PA, Lee JC, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004;304:1497-1500.
 22. Minna JD, Gazdar AF, Sprang SR, et al. Cancer: a bull's eye for targeted lung cancer therapy. *Science* 2004;304:1458-1461.
 23. Dingemans AM, Witlox MA, Stallaert RA, et al. Expression of DNA topoisomerase IIalpha and topoisomerase IIbeta genes predicts survival and response to chemotherapy in patients with small cell lung cancer. *Clin Cancer Res* 1999;5:2048-2058.
 24. Di Leo A, Isola J. Topoisomerase II alpha as a marker predicting the efficacy of anthracyclines in breast cancer: are we at the end of the beginning? *Clin Breast Cancer* 2003;4:179-186.
 25. Johnstone RW, Ruefli AA, Lowe SW. Apoptosis: a link between cancer genetics and chemotherapy. *Cell* 2002;108:153-164.
 26. Dabholkar M, Vionnet J, Bostick-Bruton F, et al. Messenger RNA levels of XPAC and ERCC1 in ovarian cancer tissue correlate with response to platinum-based chemotherapy. *J Clin Invest* 1994;94:703-708.
 27. Metzger R, Leichman CG, Danenberg KD, et al. ERCC1 mRNA levels complement thymidylate synthase mRNA levels in predicting response and survival for gastric cancer patients receiving combination cisplatin and fluorouracil chemotherapy. *J Clin Oncol* 1998;16:309-316.
 28. Yu JJ, Mu C, Lee KB, et al. A nucleotide polymorphism in ERCC1 in human ovarian cancer cell lines and tumor tissues. *Mutat Res* 1997;382:13-20.
 29. Brown JM, Wouters BG. Apoptosis, p53, and tumor cell sensitivity to anticancer agents. *Cancer Res* 1999;59:1391-1399.
 30. Yeh JJ, Hsu WH, Wang JJ, et al. Predicting chemotherapy response to paclitaxel-based therapy in advanced non-small-cell lung cancer with P-glycoprotein expression. *Respiration* 2003;70:32-35.
 31. Kawasaki M, Nakanishi Y, Kuwano K, et al. Immunohistochemically detected p53 and P-glycoprotein predict the response to chemotherapy in lung cancer. *Eur J Cancer* 1998;34:1352-1357.
 32. Hsia TC, Lin CC, Wang JJ, et al. Relationship between chemotherapy response of small cell lung cancer and P-glycoprotein or multidrug resistance-related protein expression. *Lung* 2002;180:173-179.
 33. Savaraj N, Wu CJ, Xu R, et al. Multidrug-resistant gene expression in small-cell lung cancer. *Am J Clin Oncol* 1997;20:398-403.
 34. Rosell R, Scagliotti G, Danenberg KD, et al. Transcripts in pretreatment biopsies from a three-arm randomized trial in metastatic non-small-cell lung cancer. *Oncogene* 2003;22:3548-3553.
 35. Dingemans AC, van Ark-Otte J, Span S, et al. Topoisomerase IIalpha and other drug resistance markers in advanced non-small cell lung cancer. *Lung Cancer* 2001;32:117-128.
 36. Nakanishi Y, Kawasaki M, Bai F, et al. Expression of p53 and glutathione S-transferase-pi relates to clinical drug resistance in non-small cell lung cancer. *Oncology* 1999;57:318-323.
 37. Lord RV, Brabender J, Gandara D, et al. Low ERCC1 expression correlates with prolonged survival after cisplatin plus gemcitabine chemotherapy in non-small cell lung cancer. *Clin Cancer Res* 2002;8:2286-2291.
 38. Ryu JS, Hong YC, Han HS, et al. Association between polymorphisms of ERCC1 and XPD and survival in non-small-cell lung cancer patients treated with cisplatin combination chemotherapy. *Lung Cancer* 2004;44:311-316.
 39. Camps C, Sarries C, Roig B, et al. Assessment of nucleotide excision repair XPD polymorphisms in the peripheral blood of gemcitabine/cisplatin-treated advanced non-small-cell lung cancer patients. *Clin Lung Cancer* 2003;4:237-241.
 40. Gregorc V, Ludovini V, Pistola L, et al. Relevance of p53, bcl-2 and Rb expression on resistance to cisplatin-based chemotherapy in advanced non-small cell lung cancer. *Lung Cancer* 2003;39:41-48.
 41. Rodenhuis S, Boerrigter L, Top B, et al. Mutational activation of the K-ras oncogene and the effect of chemotherapy in advanced adenocarcinoma of the lung: a prospective study. *J Clin Oncol* 1997;15:285-291.
 42. Krug LM, Miller VA, Filippa DA, et al. Bcl-2 and bax expression in advanced non-small cell lung cancer: lack of correlation with chemotherapy response or survival in patients treated with docetaxel plus vinorelbine. *Lung Cancer* 2003;39:139-143.
 43. Takayama K, Ogata K, Nakanishi Y, et al. Bcl-2 expression as a predictor of chemosensitivities and survival in small cell lung cancer. *Cancer J Sci Am* 1996;2:212.